

# Roots of Polynomials

**Spoken Tutorial Project**

**<http://spoken-tutorial.org>**

**National Mission on Education through ICT**

**<http://sakshat.ac.in>**

**Vidhya Iyer**

**IIT Bombay**

**9 June 2018**



# Learning Objectives



# Learning Objectives

**We will learn,**



# Learning Objectives

**We will learn,**

- **To plot graphs of polynomial equations**



# Learning Objectives

We will learn,

- To plot graphs of polynomial equations
- About complex numbers, real and imaginary roots



# Learning Objectives

We will learn,

- To plot graphs of polynomial equations
- About complex numbers, real and imaginary roots
- To find extrema and inflection points



# Pre-requisites



# Pre-requisites

- **GeoGebra interface**





# Pre-requisites

- **GeoGebra interface**
- **Basics of coordinate system**



# Pre-requisites

- **GeoGebra interface**
- **Basics of coordinate system**
- **Polynomials**



# Pre-requisites

- **GeoGebra interface**
- **Basics of coordinate system**
- **Polynomials**
- **If not, for relevant tutorials, please visit our website**  
**[www.spoken-tutorial.org](http://www.spoken-tutorial.org)**



# System Requirement



# System Requirement

- **Ubuntu Linux OS v 14.04**



# System Requirement

- **Ubuntu Linux OS v 14.04**
- **GeoGebra 5.0.388.0-d**



# Binomial Theorem



# Binomial Theorem

- Binomial theorem states that if  $a, b \in \mathbb{R}$ , index  $n$  is a positive integer,





# Binomial Theorem

- Binomial theorem states that if  $a, b \in \mathbb{R}$ , index  $n$  is a positive integer,
- $0 \leq r \leq n$ , then



# Binomial Theorem

- Binomial theorem states that if  $a, b \in \mathbb{R}$ , index  $n$  is a positive integer,
- $0 \leq r \leq n$ , then
- $(a + b)^n = {}^nC_0 a^n + {}^nC_1 a^{n-1} b^1 + {}^nC_2 a^{n-2} b^2 + \dots + {}^nC_r a^{n-r} b^r + \dots + {}^nC_n b^n$



# Binomial Theorem

- Binomial theorem states that if  $a, b \in \mathbb{R}$ , index  $n$  is a positive integer,
- $0 \leq r \leq n$ , then
- $(a + b)^n = {}^nC_0 a^n + {}^nC_1 a^{n-1} b^1 + {}^nC_2 a^{n-2} b^2 + \dots + {}^nC_r a^{n-r} b^r + \dots + {}^nC_n b^n$
- **Reminder,**  ${}^nC_1 = \frac{n!}{1!(n-1)!}$



# Quadratic Equations and Roots



# Quadratic Equations and Roots

- A second degree polynomial,  $y = ax^2 + bx + c$  has roots  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$  where  $\Delta = b^2 - 4ac$



# Quadratic Equations and Roots

- A second degree polynomial,  $y = ax^2 + bx + c$  has roots  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$  where  $\Delta = b^2 - 4ac$
- When  $\Delta < 0$ , roots are complex



# Quadratic Equations and Roots

- A second degree polynomial,  $y = ax^2 + bx + c$  has roots  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$  where  $\Delta = b^2 - 4ac$
- When  $\Delta < 0$ , roots are complex
- When  $\Delta = 0$ , roots are real and equal



# Quadratic Equations and Roots

- A second degree polynomial,  $y = ax^2 + bx + c$  has roots  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$  where  $\Delta = b^2 - 4ac$
- When  $\Delta < 0$ , roots are complex
- When  $\Delta = 0$ , roots are real and equal
- When  $\Delta > 0$ , roots are real and unequal





# Quadratic Equations and Roots



# Quadratic Equations and Roots

- When roots are real,  $ax^2 + bx + c = 0$  has extremum  $(x_v, y_v)$



# Quadratic Equations and Roots

- When roots are real,  $ax^2 + bx + c = 0$  has extremum  $(x_v, y_v)$
- $x_v = \frac{-b}{2a}$  and  $y_v = ax_v^2 + bx_v + c$



# Complex Numbers, XY Plane



# Complex Numbers, XY Plane

- A complex number,  $z = a + bi$  where  $a =$  real part,  $bi =$  imaginary part, and  $a$  and  $b$  are constants



# Complex Numbers, XY Plane

- **A complex number,  $z = a + bi$  where  $a =$  real part,  $bi =$  imaginary part, and  $a$  and  $b$  are constants**
- **Imaginary number,  $i = \sqrt{-1}$**



# Complex Numbers, XY Plane

- A complex number,  $z = a + bi$  where  $a =$  real part,  $bi =$  imaginary part, and  $a$  and  $b$  are constants
- Imaginary number,  $i = \sqrt{-1}$
- In the XY plane,  $a + bi$  is point  $(a, b)$



# Complex Numbers, XY Plane

- A complex number,  $z = a + bi$  where  $a =$  real part,  $bi =$  imaginary part, and  $a$  and  $b$  are constants
- Imaginary number,  $i = \sqrt{-1}$
- In the XY plane,  $a + bi$  is point  $(a, b)$
- In the complex plane, x axis = real axis, y axis = imaginary axis






# Complex Numbers, Complex Plane




# Complex Numbers, Complex Plane



- In complex plane,  $z$  is a vector with real axis co-ordinate  $a$  and imaginary axis co-ordinate  $b$




# Complex Numbers, Complex Plane



- In complex plane,  $z$  is a vector with real axis co-ordinate  $a$  and imaginary axis co-ordinate  $b$
- Length of vector  $z = |z| = r$



# Complex Numbers, Complex Plane



- In complex plane,  $z$  is a vector with real axis co-ordinate  $a$  and imaginary axis co-ordinate  $b$
- Length of vector  $z = |z| = r$
- $r = \sqrt{a^2 + b^2}$  (Pythagoras' theorem)



# Point of Inflection



# Point of Inflection

**Point of inflection (Pol) on a curve is the point where curve changes direction.**

- **To find co-ordinates of Pol  $(x, y)$ , we equate 2nd derivative of given function to 0**



# Point of Inflection

**Point of inflection (Pol) on a curve is the point where curve changes direction.**

- **To find co-ordinates of Pol  $(x, y)$ , we equate 2nd derivative of given function to 0**
- **Solve to get  $x$  ( $x$  co-ordinate of Pol)**



# Point of Inflection

**Point of inflection (Pol) on a curve is the point where curve changes direction.**

- To find co-ordinates of Pol  $(x, y)$ , we equate 2nd derivative of given function to 0
- Solve to get  $x$  ( $x$  co-ordinate of Pol)
- Substitute this  $x$  in original function to get  $y$  co-ordinate





# Summary

**In this tutorial, we have learnt to,**

- **Plot graphs of polynomial functions using CAS view and input bar**
- **Find real roots, extrema and inflection point(s)**

**Complex roots will be covered in another tutorial**



# Assignment



# Assignment

- $d(x) = x^2 - 6x + 5$



# Assignment

- $d(x) = x^2 - 6x + 5$
- $e(x) = 3x^3 - 2x^2 + 0.2x - 1$



# Assignment

- $d(x) = x^2 - 6x + 5$
- $e(x) = 3x^3 - 2x^2 + 0.2x - 1$
- $f(x) = 2x^4 - x^3 + 3x^2$



# Assignment

- $d(x) = x^2 - 6x + 5$
- $e(x) = 3x^3 - 2x^2 + 0.2x - 1$
- $f(x) = 2x^4 - x^3 + 3x^2$
- $g(x) = x^5 - 7x^4 + 9x^3 + 23x^2 - 50x + 24$



# Assignment

- $d(x) = x^2 - 6x + 5$
- $e(x) = 3x^3 - 2x^2 + 0.2x - 1$
- $f(x) = 2x^4 - x^3 + 3x^2$
- $g(x) = x^5 - 7x^4 + 9x^3 + 23x^2 - 50x + 24$
- $h(x) = (4x + 3)/(x - 1)$



# Assignment

- $d(x) = x^2 - 6x + 5$
- $e(x) = 3x^3 - 2x^2 + 0.2x - 1$
- $f(x) = 2x^4 - x^3 + 3x^2$
- $g(x) = x^5 - 7x^4 + 9x^3 + 23x^2 - 50x + 24$
- $h(x) = (4x + 3)/(x - 1)$
- $i(x) = (3x^2 - 2x - 1)/(2x^2 + 3x - 2)$





# About the Spoken Tutorial Project

- Watch the video available at [http://spoken-tutorial.org/What\\_is\\_a\\_Spoken\\_Tutorial](http://spoken-tutorial.org/What_is_a_Spoken_Tutorial)
- It summarizes the Spoken Tutorial project
- If you do not have good bandwidth, you can download and watch it



# Spoken Tutorial Workshops

## The Spoken Tutorial Project Team

- Conducts workshops using spoken tutorials
- Gives certificates to those who pass an online test
- For more details, please write to [contact@spoken-tutorial.org](mailto:contact@spoken-tutorial.org)



# Forum for specific questions

- Do you have questions in **THIS Spoken Tutorial?**
- Please visit <http://forums.spoken-tutorial.org>
- Choose the minute and second where you have the question
- Explain your question briefly
- Someone from our team will answer



# Acknowledgements

- Spoken Tutorial Project is a part of the Talk to a Teacher project
- It is supported by the National Mission on Education through ICT, MHRD, Government of India
- More information on this Mission is available at

<http://spoken-tutorial.org /NMEICT-Intro>

